

CONTRIBUTION TO THE LICHEN FLORA OF BRAZIL XI.  
Lichens from Santa Maria, Rio Grande do Sul State.

HECTOR S. OSORIO \* and MARIANA FLEIG \*\*.

\*Departamento de Botánica, Museo Nacional de  
Historia Natural, Casilla de Correo 399,  
Montevideo, URUGUAY.

\*\*Departamento de Botanica, Instituto de Bio-  
ciencias, Universidade Federal do Rio Grande  
do Sul, PORTO ALEGRE, RS BRASIL.

The Municipality of Santa Maria, located in the central part of Rio Grande do Sul State, is the westernmost region of the Central Lowlands (Depressao Central, Rambo 1956).

The present knowledge we possess of its lichen flora is due exclusively to the collections made by G. A. Malme during the First Regnell Expedition (Malme 1897). During the Second Regnell Expedition (Malme 1904) this Municipality was not visited by this author.

Apart from the works dealing with the collections made during the First Regnell Expedition the authors were unable to find any other information concerning the lichen flora of this Municipality.

The present paper is based on a collection made by the junior author in the place named "Parada Link" which is a small railway station located 25 km NW from Santa Maria City.

All the lichen species here enumerated were collected growing on Melia azedarach and are preserved at ICN. The collection numbers belong to the junior author numbering system.

Buellia callispora (Nyl.) Steiner.

703/28b. New to Rio Grande do Sul.

Caloplaca granularis (Müll. Arg.) C. Sambo.

703/3b, 703/26b. In Brazil this species is at present known only from Rio Grande do Sul State: two collections from the vicinity of Encruzilhada do Sul (Osorio & Homrich 1978) and two another collections from the Municipality of Montenegro (Osorio, Aguiar & Citadini 1980).

Dirinaria appianata (Fée) Awasthi.

703/3a.

Glyphis cicatricosa (Ach.) Vain. f. confluens (Zenk.)  
703/20. Zahlbr.

Redinger (1934) reported three collections (including one of the forma intermedia) from the whole

State of Rio Grande do Sul. Field observations made by the authors in several localities proved that at present this species is widely distributed through the State. It was observed growing indistinctly on indigenous or introduced trees.

Graphina nylanderiana Zahlbr.

703/23.

Graphis lineola Ach.

703/22.

Haematomma similis Bagl.

703/16, 703/32. Malme (1940) who reported this species as Haematomma puniceum var. subinnatum do not quote any collection from Rio Grande do Sul. Rogers (1982) in the lectotypification of this variety mentions erroneously the locality of Colonia Risso from Uruguay instead of Paraguay.

Heterodermia diademata (Tayl.) Awasthi.

703/1, 703/9.

Lecidea russula Ach.

703/19.

Ochrolechia subpallescens Vers.

703/26a, 703/30. Formerly known in the State from only one collection made near Encruzilhada do Sul (Osorio & Homrich 1978).

Parmelina lindmanii (Lyngé) Hale.

703/5, 703/6.

Parmotrema austrosinense (Zahlbr.) Hale.

703/8.

Parmotrema praesorediosum (Nyl.) Hale.

703/7.

Pertusaria flavens Nyl.

703/29a.

Pertusaria pulchella Malme.

703/18.

Phaeographina arechavaletae Müll. Arg.

703/24. Formerly known in Brazil from only one collection made near Encruzilhada do Sul in Rio Grande do Sul State (Osorio & Homrich 1978).

Phaeographina caesiopruinosa (Fée) Müll. Arg.

703/21.

Phaeographis lobata (Eschw.) Müll. Arg.

703/28a. Formerly known from two localities placed in the eastern region of the State (Osorio & Fleig 1982, Osorio, Homrich & Fleig 1982).

Phaeographis medusiformis (Kremplh.) Müll. Arg.

703/17.

Physcia aipolia (Ehrh.) Hampe.

703/15.

Physcia crispa Nyl.

703/2.

Pseudoparmelia carneopruinata (Zahlbr.) Hale.

703/13.

Pseudoparmelia exornata (Zahlbr.) Hale.

703/14.

Pseudoparmelia texana (Tuck.) Hale.

703/4.

Ramalina celastri (Spreng.) Krog & Swinsc.

703/11.

Sphinctrina depressa H. Magn.

703/29c. New to Brazil. Formerly known only from two localities in Uruguay (Magnusson 1950. Osorio 1979).

#### SUMMARY.

Twenty six lichen species collected in the Municipality of Santa Maria are listed. Buellia callispora and Haematomma similis are added to the known flora of Rio Grande do Sul State. Sphinctrina depressa is recorded for Brazil for first time.

#### LITERATURE CITED.

- MAGNUSSON, A. H. 1950. Lichens from Uruguay. *Meddelanden för Göteborgs Botaniska Trädgård* 18: 213-237.
- MALME, G. O. 1904. Reseberättelse afgifven at Regnellske stipendiats doktor G.O. Malme för aren 1901-1903. *Kunglige Svenska Vetenskaps-Akademiens Årsbok* 1904: 105-115.
- MALME, G. O. 1940. Lichenes nonnulli in *Expeditione Regnelliana prima collecti*. *Arkiv för Botanik* 29A (6): 1-35.
- OSORIO, H. S. 1979. Contribution to the lichen flora of Uruguay XII. Lichens from Nueva Palmira, Colonia Department. *Phytologia* 43 (3): 289-292.
- OSORIO, H. S., L. W. AGUIAR & V. CITADINI. 1980. Contribution to the lichen flora of Brazil. VII. Lichens from Montenegro and Triunfo, Rio Grande do Sul State. *Comunicaciones botánicas del Museo de Historia Natural de Montevideo* 4(62): 1-8.
- OSORIO, H. S. & M. FLEIG. 1982. Contribution to the lichen flora of Brazil IX. Lichens from the Municipality of Torres, Rio Grande do Sul State. *Mycotaxon* 14(1): 347-350.
- OSORIO, H. S. & M. H. HOMRICH. 1978. Contribution to the lichen flora of Brazil IV. Lichens from Southern Rio Grande do Sul. *The Bryologist* 81(3): 452-454.
- OSORIO, H. S., M. H. HOMRICH & M. FLEIG. 1982. Contribution to the lichen flora of Brazil X. Lichens from Guaiba, Rio Grande do Sul State. *Phytologia* 51(7): 479-484.
- RAMBO, J. 1956. A Fisionomia do Rio Grande do Sul. Jesuitas no Sul do Brasil vol. VI: 1-456. 2a. edic. Edit! Livraria Selbach, Porto Alegre, RS, Brasil.
- REDINGER, K. 1934. Die Graphidineen der ersten Regnelli'schen Expedition nach Brasilien 1892-1894. I. Glyphis, Medusulina und Sarcographa. *Arkiv för Botanik* 25A (13): 1-20.
- ROGERS, R. W. 1982. The corticolous species of Haematomma in Australia. *Lichenologist* 14 (2): 115-129.